




Pyrethroid



Capsule Suspension



Food Handling

# PRESCRIPTION TREATMENT<sup>®</sup> brand **CY-KICK<sup>®</sup> CS**

## Controlled Release Cyfluthrin

For Use by Pest Control Operators or Commercial Applicators.

### KILLS:

Ants, Bed Bugs, Bees, Boxelder Bugs, Cadelles, Carpenter Ants, Carpenter Bees, Carpet Beetles, Centipedes, Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Cockroaches, Crickets, Darkling Beetles, Dermestids, Drugstore Beetles, Earwigs, Elm-leaf Beetles, Fire Ants, Firebrats, Fleas, Flies, Flour Beetles, Grain Weevils, Grasshoppers, Hide Beetles, Hornets, Indian Meal Moths, Larder Beetles, Lesser Grain Borers, Lesser Mealworms, Loopers, Mediterranean Flour Moths, Merchant Grain Beetles, Mealworms, Millipedes, Mosquitoes, Psocids, Rice Weevils, Sawtoothed Grain Beetles, Scorpions, Silverfish, Sowbugs, Spiders, Springtails, Termites, Ticks, Yellow Jackets, Warehouse Moths, Wasps, Wood Wasps, Wood Infesting Borers and Beetles.

### FOR USE IN AND AROUND:

Apartments, Calf Hutches, Calving Pens and Parlors, Campgrounds, Empty Chicken Houses, Dairy Areas, Dog Kennels, Food Storage Areas, Grain Mills, Granaries, Hog Barns, Homes, Horse Barns, Hospitals, Hotels, Meat Packing Plants, Food Processing Plants, Milkrooms, Motels, Nursing Homes, Rabbit Hutches, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes†), Utilities, Warehouses, Commercial and Industrial Buildings.

† Do not use in aircraft cabins.

### ACTIVE INGREDIENT:

Cyfluthrin ..... 6.00%

OTHER INGREDIENTS: ..... 94.00%

TOTAL: 100.00%

EPA Reg. No. 499-304

## KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements.

### PRECAUTIONARY STATEMENTS

#### FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment.

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

#### GENERAL INFORMATION

Cy-Kick CS Controlled Release Cyfluthrin is a flowable microencapsulated concentrate containing 0.5 lb. of cyfluthrin per gallon. Cy-Kick CS may be used on any surface which will not be damaged or stained by water. Heavy applications may leave a visible deposit. On some surfaces this deposit can be removed with a damp cloth or sponge.

In situations where conventional application methods have not or are not likely to provide adequate coverage, foam generating equipment or similar machines can be used to provide a continuous coverage. Use drier foam inside wall voids to avoid any water damage or electrical short circuits. Refer to the section entitled "Foam Application Use Directions".

Do not apply this product through any type of irrigation system.

Do not apply water-based sprays of this product to conduits, motor housing, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not spray where electrical short circuits might result, such as wall outlets, conduits, etc.

Do not apply in aircraft cabins.

Do not apply to humans, their clothing or bedding. Do not allow children or pets to contact treated surfaces until spray has dried. Do not contaminate food or use on household pets. Cover fish tanks prior to application. Do not apply this product in patient rooms while occupied or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms when in use.

Avoid contamination of food, feedstuffs, or water supply. Do not contaminate food preparation surfaces, kitchen utensils, dishes, or feed storage containers. Any food/feed contact surfaces and cooking utensils in the treatment area should be covered during treatment or thoroughly cleaned before using.

### MIXING INSTRUCTIONS

Thoroughly clean spray equipment before using Cy-Kick CS. Shake concentrate well before diluting. When diluting, first add approximately one-half of the water to the spray tank and then add the proper amount of Cy-Kick CS. When emptying the bottles, triple rinse with water, shake well and add the rinseate to the spray tank. Add the rest of the water and agitate the sprayer thoroughly. Agitate sprayer occasionally during use to ensure even mixture. Shake or re-agitate sprayer if spraying is interrupted or if dilution is left in sprayer overnight. If spray screens are used, they should be 50 mesh or larger.

#### SPRAY DILUTION CHART

| Amount of Finished Spray | Spray Concentration |              |
|--------------------------|---------------------|--------------|
|                          | 0.05% Spray         | 0.10% Spray  |
| 1 gallon                 | 1.0 fl. oz.         | 2.0 fl. oz.  |
| 3 gallons                | 3.0 fl. oz.         | 6.0 fl. oz.  |
| 5 gallons                | 5.0 fl. oz.         | 10.0 fl. oz. |
| 10 gallons               | 10.0 fl. oz.        | 20.0 fl. oz. |

The higher rate should be used for clean-outs, for longer residual control or for high insect infestations.

#### VOID / CRACK & CREVICE<sup>®</sup> APPLICATION

To treat insects harbored in voids and in cracks and crevices, apply with equipment designed for void applications. Follow the manufacturer's recommendations for proper equipment set up and operating pressures. Place the tip of the applicator at or into the crack or crevice to be treated. Apply product in short bursts while keeping tip in position. Allow the air stream to push insecticide deep into voids. Application should be made in such a manner to limit dripping and run-off on structural surfaces and plants.

#### INDOOR TREATMENTS

Dilute Cy-Kick CS to 0.05% or 0.10% with water using the spray dilution chart.



**WHITMIRE MICRO-GEN**  
RESEARCH LABORATORIES, INC.  
800-777-8570

**NOTE:** This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.



# CY-KICK CS® Controlled Release Cyfluthrin

Cy-Kick CS should be used as a Crack & Crevice residual product where insects may be hiding, living and breeding and/or for selected spot application.

When using in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed, refer to the section entitled "Food/Feed Areas of Food Handling Establishments" for use directions. Non-food/feed areas are areas such as garbage rooms, restrooms, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

**CENTIPEDES, COCKROACHES, CRICKETS, DERMESTIDS, EARWIGS, FIREBRATS, MILLIPEDES, PSOCIDS, SILVERFISH, SOWBUGS, SPIDERS:** Make spot treatment and/or Crack & Crevice injections into hiding places such as cracks and crevices, void areas, moist areas, openings around pipes and sinks and under refrigerators. Repeat as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**FLIES:** Make spot applications of Cy-Kick CS on surfaces where flies rest in the dark at night or frequent during the day, such as window frames, screens, trash cans, ceilings and upper wall surfaces.

**CARPET BEETLES (DERMESTIDS):** Treat along edges of carpet, under carpet edge, furniture, throw rugs and as a Crack & Crevice application in closets and other shelving where infested items are stored. Do not treat clothing with this product. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**BED BUGS:** Allow treated articles to dry thoroughly before use. Take bed apart. Treat joints and channels if hollow, such as square or round tubing. See that the interior framework is treated. Treat mattresses, especially tufts, folds and edges. Cover mattresses with outer sheets before use. Do not apply to other bedding articles. Also treat other areas where bed bugs may be harboring.

**TICKS:** Apply as a spot and/or Crack & Crevice treatment in infested and egg laying areas such as pet resting quarters, nearby cracks and crevices, behind baseboards, door and window frames. Replace pet's old bedding with clean fresh bedding after treatment. Do not treat pets with this product. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**ANTS:** Inject cracks, crevices and voids where ants may be traveling and/or harboring, including but not limited to, around doors, window frames, and other structural cracks, crevices and voids. Spot treatment of trails and feeding areas may also be made.

**STORED PRODUCT PESTS (EXPOSED ADULT AND IMMATURE STAGES):** Before storing products, treat warehouses, production facilities, storage areas, rail cars, truck beds and other areas where products are stored with Cy-Kick CS. Apply as a spot and/or Crack & Crevice treatment to cracks, crevices and surfaces where pests have been seen or may have harborage. Cy-Kick CS may be used to treat grain storage facilities and other listed areas for stored product pest control. Cleaning areas prior to use of this product will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

**FOAM APPLICATION USE DIRECTIONS:** Cy-Kick CS can be converted to a foam and the foam used to treat structural voids to control insect and arthropod pests in walls, under slabs or in other void areas. Apply as a dry foam (approximately 25:1). Fill void with foam so that all surfaces are contacted. Mix end-use dilution of Cy-Kick CS with manufacturer's recommended volume of foaming agent. Use the recommended rate suspension converted to a foam to achieve the desired foam expansion ratios. Verify before treatment that the foaming agent is compatible with Cy-Kick CS.

### FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

Cy-Kick CS applications are permitted in food/feed areas of food/feed handling establishments as a General Surface Application, Spot or Crack & Crevice Treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Food/feed areas are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.

**GENERAL SURFACE APPLICATION:** Do not apply as a general surface application in the food/feed handling areas of the facility when the facility is in operation or foods are exposed. Do not apply directly to food products. Cover or remove all food processing and/or handling equipment during application. After application in food processing plants, bakeries, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food will contact. Clean food handling or processing equipment and thoroughly rinse with clean, fresh water.

**SPOT AND CRACK & CREVICE APPLICATION:** Spot or Crack & Crevice applications may be made while the facility is in operation; however, food should be covered or removed from area being treated. Do not apply directly to food or food-handling surfaces. May be used in federally inspected meat and poultry plants.

### INDOOR AND OUTDOOR TREATMENTS

Select one of the following dilutions, 0.05% or 0.10%, for insects listed below unless otherwise noted.

**HIBERNATING STAGES OF BOXELDER BUGS, ELM-LEAF BEETLES, CLUSTER FLIES AND CLOVER MITES:** Inject cracks, crevices and/or voids where these insects may be harboring or hibernating. Infested areas include, but are not limited to, cracks and crevices in weep holes, wall voids, around window and door frames, and under siding. Treat wall area around entry points on exterior where insects congregate. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**CARPENTER ANTS:** Make spot and/or Crack & Crevice applications to any trails, inside or outside around doors and windows, and other places where ants enter premises. Areas which may be treated include, but are not limited to, atriiums, crawl spaces, ceilings, garages, soffits, wall voids, weep holes, and other void areas. Where possible, treat ant nest or infested wood directly. Repeat application as necessary, but not more than twice per week. Treatments must be at least 3 days apart.

**BEES, WASPS AND YELLOW JACKETS:** As a preventative or remedial treatment to help prevent these insects from building nests or workers from congregating on structures, follow perimeter treatment directions under outdoor section of label. Pay particular attention to soffit and eave areas.

For treatment of a yellow jacket or wasp nest, liberally treat the nest, inside the nest for yellow jackets and the surrounding area. For treatment of nests inside wall voids, inject into the nest sufficient quantity to treat the complete nest. More than one point of application may be required. Most applications should require no more than half a gallon of diluted material. Pay particular attention to what the water may contact such as electrical wiring. Do not use where there is a shock hazard.

CY-KICK CS DOES NOT PROVIDE INSTANT KNOCKDOWN. THE BEST TIME TO TREAT IS LATE IN THE DAY. COLONY SHOULD BE DEAD IN 2-3 DAYS. DEPENDING ON THE SITUATION, PROTECTIVE EQUIPMENT FOR THE APPLICATOR MAY BE REQUIRED.

### WOOD DESTROYING INSECTS

**LOCALIZED TREATMENT OF WOOD-INFESTING INSECTS - Beetles and Borers (such as the following families):** Anobiidae, Bostriichidae, Buprestidae, Cerambycidae, Lyctidae, Oedemeridae, Sesiidae and Termitidae. Locate suspected infestation and/or insect tunnels and cavities. Drill pilot holes into insect tunnels, gallery and inaccessible closed voids. Drill holes 8 to 10 inches apart to ensure thorough coverage. Inject 5 to 10 seconds per hole. Do not contact electrical wiring, plumbing, etc. Use the 0.05% or 0.10% rate.

**TERMITES:** To inhibit termite infestations, treat exposed tubing, cracks or cold joints in concrete or other structural interstices. Treatment of exposed surfaces should be done to cover thoroughly but excess run-off should be avoided. This type of application is intended to be a supplemental treatment for control of drywood and above ground subterranean termites and other wood-infesting insects. It is not intended to be a substitute for soil treatment or for extensive infestation of drywood or other wood-infesting insects. Use a 0.125% rate, 2.5 fl. oz. of concentrate per gallon of water. To treat for severe infestations or for treatments under slabs, in basements and buildings with insufficient crawlspace, use a registered termiticide.

### OUTDOOR TREATMENTS

**KILLS:** Ants, Boxelder Bugs, Carpenter Ants, Carpenter Bees, Carpet Beetles, Centipedes, Clover Mites, Cluster Flies, Cockroaches, Crickets, Dermestids, Earwigs, Elmleaf Beetles, Fleas, Flies, Grasshoppers, Millipedes, Mosquitoes, Psocids, Scorpions, Silverfish, Sowbugs, Spiders and Ticks.

Cy-Kick CS can be used as a residual treatment on outdoor surfaces and/or as a crack, crevice and void treatment of buildings, porches, window frames, eaves, patios, garages, garbage dumps and other areas where pests congregate or enter premises. Repeat treatment as necessary to maintain effectiveness, but not more than twice per week. Treatments must be at least 3 days apart. Use the 0.05% or 0.10% rate.

Cy-Kick CS can be used as a residual treatment to control pests in outdoor void areas such as meter boxes and manholes. Repeat treatment as necessary, but not more than twice per week. Treatments must be at least 3 days apart. Use the 0.05% or 0.10% rate.

**PERIMETER TREATMENT:** To help prevent infestations of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure. Cy-Kick CS may be applied in compressed air, backpack or power spray equipment. An application rate of 0.5 to 1.0 fluid ounce of Cy-Kick CS per 1000 sq. ft. is desirable. Apply in sufficient water to adequately cover the area being treated. Application should be made in such a manner to limit dripping and run-off on structural surfaces and plants. Alternate directions are to use 4 to 8 fl. oz. of product per 50 gallons of water.

**FLIES:** Treat outside surfaces of screens, doors, window frames or wherever flies enter. Also treat surfaces on porches, in garages, and other places where insects light or congregate. Repeat as necessary. Treat normal fly resting areas such as ceilings, upper corners of the area, around windows and doors, and

other surfaces. Treat inside and outside surfaces of dumpsters and other trash holding containers. Treat trash storage areas. Treat so as to lightly moisten the target surface. For best results, begin treatments at the start of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

**MOSQUITO CONTROL OUTDOORS:** To control adult mosquitoes, use handheld, back-pack or truck mounted sprayers only. Not for use with truck mounted cold aerosol ULV sprayers, hand-held cold aerosol ULV sprayers or thermal fogging devices. Do not apply by aircraft. Do not apply through any kind of irrigation and/or lawn sprinkler systems. Use up to 150 gallons of diluted spray per acre. Treat shrubbery, vegetation and other areas where mosquitoes may rest. Shrubby vegetation and other areas where mosquitoes, ponds and shorelines may be treated if applied as a wet spray. Direct application of this product to any body of water is prohibited.

**CARPENTER BEES:** Locate entrance holes. Treat galleries through entrance holes. Seal bee entrances after application.

**SCORPIONS:** Remove accumulations of lumber, firewood, and other materials which serve as harborage sites. Before stacking firewood or lumber, apply this product as a residual spray to surfaces immediately below such materials. Band treatment may also be helpful in reducing pests immigrating from surrounding areas.

**CRAWLING INSECTS - including but not limited to Ants, Cockroaches, Crickets, Earwigs, Sowbugs, and Scorpions:** For control of infestations of these insects in outdoor areas which are landscaped, planted, mulched or unmaintained. For ants, treat mounds and runways. Repeat applications as infestation warrants and as reinfestations occur. Apply Cy-Kick CS to cracks and crevices where insects may enter premises. Areas to be treated include, but are not limited to, weep holes, under siding, wall voids, soffits, around attic vents, cracks and crevices, openings around windows, doors and pipes. Treatment of infested fences and tree holes is permitted.

**FLEAS:** Treat infested areas. Cy-Kick CS will kill fleas on contact but does not provide residual control.

### USE IN AND AROUND LIVESTOCK HOUSING

MAY BE USED IN AND ON OUTSIDE SURFACES OF OTHER BUILDINGS. Do not apply to lactating or food animals. Do not contaminate milk, food or drinking water. Remove all animals prior to making application to housing. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until surfaces are dry.

**HOUSE FLIES AND STABLE FLIES:** For residual control of these pests on outdoor surfaces and in animal areas such as dog kennels, hog barns, horse barns, empty chicken houses, rabbit hutches, dairy areas, milkrooms, calf hutches, calving pens and parlors. Treat normal fly resting areas such as ceilings, screens, upper corners of the area, around windows and doors, light fixtures and other surfaces known to attract flies. Apply as a coarse wet spray. Using a mixture of 1 fl. oz. of Cy-Kick CS per gallon of water; apply at a rate of 1 gallon diluted product per 1,000 sq. ft. of surface. For best results initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity. Treat no more than twice per week and at least 3 days apart.

**PERIMETER TREATMENT:** For control of other pests and to help prevent infestations of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to the building. Also treat the building foundation and sides where pests are active and may find entrance. Treat other insect entry points such as doors, windows, under siding and eaves of structure. Cy-Kick CS may be applied in compressed air, backpack or power spray equipment. An application rate of 1.0 fluid ounces of Cy-Kick CS per 1000 sq. ft. is desirable. Apply in sufficient water to adequately cover the area being treated. Application should be made in such a manner to limit dripping and run-off on structural surfaces and plants.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides or fertilizers. Do not store above 122°F for extended periods of time. Preferred storage is above 32°F. Storage below 32°F may result in solidification. If warmed to above 32°F, product will return to original form. Freezing does not adversely affect this product. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL (1 gal.):** Empty container by using the product in accordance with the label directions. Do not reuse container! Place empty container in trash or offer for recycling if available. If container is partly filled, call your local solid waste agency or call 1-800-CLEANUP for disposal instructions. NEVER PLACE UNUSED PRODUCT DOWN ANY INDOOR OR OUTDOOR DRAIN!

**CONTAINER DISPOSAL (Greater than 1 gal.):** Empty container by using the product in accordance with the label directions. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### A Prescription Treatment® brand insecticide from:

Whitmire Micro-Gen Research Laboratories, Inc.  
3568 Tree Court Industrial Blvd.  
St. Louis MO 63122-6682

© 2005 Whitmire Micro-Gen Research Laboratories, Inc.

**NOTE:** This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.