



Kenya Pyrethrum



Synergized



System Equipment



Food Handling

Prescription Treatment® brand

P.I.®

Contact Insecticide

Formula 1

KILLS:

Angoumois Grain Moths, Ants, Bed Bugs, Booklice, Carpet Beetles, Centipedes, Chocolate Moths, Cigarette Beetles, Clothes Moths, Clover Mites, Cluster Flies, Cockroaches, Confused Flour Beetles, Crickets, Drug Store Beetles, Earwigs, Fleas, Flies, Fruit Flies, Gnats, Grain Mites, Granary Weevils, Horn Flies, House Flies, Indian Meal Moths, Mediterranean Flour Moths, Millipedes, Mosquitoes, Mud Daubers, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Silverfish, Small Flying Moths, Sowbugs, Spiders, Stable Flies, Ticks and Wasps.

FOR USE IN AND AROUND:

Apartments, Campgrounds, Food Storage Areas, Homes, Hospitals†, Hotels, Motels, Nursing Homes†, Resorts, Restaurants†, and other Food Handling Establishments†, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks and Aircraft – Cargo Areas Only), Utilities, Warehouses and other Commercial and Industrial Buildings.

† See special instructions for these sites under Directions for Use.

ACTIVE INGREDIENTS:

Pyrethrins, a botanical insecticide05%
Piperonyl Butoxide, Technical	4.0%

OTHER INGREDIENTS:95.5%
---------------------------------	--------

Contains petroleum distillate.

TOTAL: 100.0%

EPA Reg. No. 499-444

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 1-800-225-3320 for emergency medical treatment.

NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Extremely Flammable. Contents under pressure. Keep away from fire, sparks and heated surfaces. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

- Use only with the supplied actuator and injection tubes or other Whitmire Micro-Gen equipment.
- Do not spray on plastic, painted or varnished

surfaces or directly into any electronic equipment such as radios, televisions, computers, etc.

- Aerosols work best when can temperature is above 60° F. If can temperature is below 60° F, store at room temperature until a temperature above 60° F is reached.

CRACK & CREVICE® TREATMENT: Inject P.I.® into cracks and crevices or void spaces where insects may be harboring, living and breeding. Place injector tip into cracks, crevices, holes and other small openings. Apply product for 1 second. For light infestations, move injector tip along cracks while treating at the rate of 3 linear feet per second. For heavy infestations, move injector tip along at 1 linear foot per second. For treatment of inaccessible voids (including wall and equipment voids), calculate the void's cubic area and treat at the rate of 5 to 10 seconds per 3 cu. ft. Several holes may be required in long-running voids.

SPACE TREATMENT: Calculate the cubic footage to be treated and apply at the dosage rate specified below for the pest(s) to be treated. Ventilate treated areas before reoccupying.

INDOOR TREATMENTS

CLUSTER FLIES, FLIES, FRUIT FLIES, GNATS, HORN FLIES, HOUSE FLIES, MOSQUITOES, SMALL FLYING MOTHS AND STABLE FLIES:



WHITMIRE MICRO-GEN
RESEARCH LABORATORIES, INC.
800-777-8570

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.



P.I.® Contact Insecticide Formula 1

Apply as a space treatment at a rate of 1 to 3 seconds per 1,000 cu. ft. Close all windows and doors. Direct fog upward and disperse in all locations. Keep area closed for 15 minutes. Open and ventilate before reoccupying.

CLUSTER FLIES, CLOVER MITES, WASPS AND BEES INSIDE STRUCTURAL VOIDS: For treatment of these pests, locate all insect entrances and close all but one opening. Determine size of void. Inject 5 to 10 seconds per 3 cu. ft. of void into remaining opening. For best results, treat at dusk or later.

ANTS, BOOKLICE, CENTIPEDES, CLOVER MITES, CRICKETS, MILLIPEDES, ROACHES, SILVERFISH, SOWBUGS AND SPIDERS: Apply as a Crack & Crevice treatment into cracks and crevices in all hiding places (such as behind baseboards, sinks, cabinets, meter boxes, door frames and windows) or as a space treatment at a rate of 20 seconds per 1,000 cu. ft. For a space treatment, open cabinets and doors in area to be treated, turn off air conditioners and fans and close doors and windows before treating. Disperse toward area suspected of harboring the greatest insect infestations. Disperse in all locations contacting as many insects as possible. Keep area closed for 15 minutes. Open and ventilate the treated area before reoccupying.

For treatment of void areas such as attics, false ceilings and crawl spaces, calculate the volume of the void and treat at a rate of 20 seconds per 1,000 cu. ft.

BED BUGS: Take bed apart. Treat joints and channels if hollow, such as square or round tubing, and see that the interior framework is treated. Treat mattresses, especially tufts, folds and edges. Also treat other areas where bed bugs may be harboring. Do not use in patient rooms in hospitals and nursing homes for treatment of bed bugs.

CARPET BEETLES (IN RESIDENCES): Hold product 36" above floor and direct spray toward floor and lower walls treating at a rate of 10 seconds per 100 sq. ft. Also apply as a Crack & Crevice treatment to cracks and crevices near the source of the infestation.

CLOTHES MOTHS: Locate source of infestation. Apply as a Crack & Crevice treatment to nearby cracks and crevices and perform a space treatment of the infested area at a rate of 5 to 10 seconds per 1,000 cu. ft.

To protect woolens and other keratin containing materials, brush clean then air out. Before treating entire article, treat a small hidden area of fabric to determine if staining will occur. Treat thoroughly by holding dispenser at least 18" from garment. Treated articles should be put into a darkened closet or other darkened storage such as cedar chests, storage

drawers or sealable plastic fabric bags. For best protections, clean and treat inside of closets or storage areas by treating 1 second per 10 sq. ft. before putting treated materials away. Treatment should be repeated every 6 months or as needed. Dry clean treated clothes before wearing.

CARPET BEETLES, CIGARETTE BEETLES, CONFUSED FLOUR BEETLES, DRUG STORE BEETLES, GRAIN MITES, GRANARY WEEVILS, RED FLOUR BEETLES, RICE WEEVILS AND SAWTOOTHED GRAIN BEETLES (FOR EXPOSED ADULT AND LARVAL STAGES): Apply as a Crack & Crevice treatment into cracks and crevices of pantries, cabinets, food processing and handling equipment and other places where insects harbor or as a space treatment at a rate of 20 seconds per 1,000 cu. ft. Also apply around cartons, containers and other areas where these insects tend to congregate. Contact as many insects as possible. Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

ANGOUMOIS GRAIN MOTHS, CHOCOLATE MOTHS, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS (FOR EXPOSED ADULT AND LARVAL STAGES): Apply as a space treatment at a rate of 5 to 10 seconds for each 1,000 cu. ft. Close all doors and windows. Direct into all parts of room, especially around stored product containers, pallets and darkened areas. Keep area closed for 15 minutes following treatment. Open and ventilate before reoccupying. Repeat treatment weekly if necessary. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

FLEAS IN BUILDINGS: Hold product 36" above floor and direct spray toward floor and lower walls treating at a rate of 10 seconds per 100 sq. ft. Make sure that all floor, sofa and chair surfaces are contacted. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days if necessary.

TICKS: Holding container 36" above the floor, direct spray toward pet beds and resting quarters treating at a rate of 10 seconds per 100 sq. ft. Also apply as a Crack & Crevice treatment into adjacent cracks and crevices such as behind baseboards and edges of carpet and floor covers. Also treat higher cracks such as upper door jams and window framing, behind pictures and other areas where females may crawl to lay eggs.

IN HOSPITALS AND NURSING HOMES
Apply according to specific directions for pest(s) to be treated. Patients should be removed from room prior

to treatment. Room should be ventilated for 2 hours after spraying. Do not return patients to room until after ventilation.

FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

Apply according to specific directions for pest(s) to be treated. When applying as a space treatment in food handling areas, the areas of the operation being treated should be shut down. In food processing areas, all utensils, shelving, etc., where food will be handled should be covered or removed before treatment or thoroughly washed with an effective cleaning compound followed by a potable water rinse prior to use.

OUTDOOR TREATMENTS GROUND APPLICATION (FLIES, GNATS AND MOSQUITOES): Treat open areas near buildings and in campgrounds. Apply at a rate of 5 to 15 seconds per 1,000 sq. ft. Apply in wide swaths across area to be treated. Allow treatment to penetrate dense foliage. Best results are obtained when wind speed is 5 MPH or less.

FLIES, CLUSTER FLIES, CLOVER MITES, MUD DAUBERS AND WASPS: Apply mist directly on insects and their nests from approximately 18". Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Disperse into hiding and breeding places contacting as many insects as possible

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool area away from heat or open flame.

PESTICIDE DISPOSAL: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not puncture or incinerate! Empty container by using the product in accordance with the label directions. Offer empty container for recycling, if available, or place in trash if allowed by state and local regulations. If container is partly full, contact your local solid waste agency or call 1-800-CLEANUP for disposal instructions.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.



A Prescription Treatment® brand insecticide from:
Whitmire Micro-Gen Research Laboratories, Inc.
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
© 2005 Whitmire Micro-Gen Research Laboratories, Inc.

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.